

Exceptional and non-exceptional contributions to the radiative π decay*

V. Mateu

Departament de Física Teòrica-IFC, Universitat de València-CSIC,

**E-mail: mateu@ific.uv.es*

We have studied the spin-one resonance dominated form factors governing the radiative decay of the π , within the framework of resonance chiral theory. We obtain predictions for their value at zero momentum transfer and also a description of their q^2 dependence. We also fit the possible new physics tensor coupling.

Keywords: Chiral lagrangians; $1/N_C$ expansions; Proceedings.

1. Main results and conclusions

For many years the radiative π decay (RPD) process ($\pi \rightarrow e\nu\gamma$) has been an open window for the speculation about new physics (NP).¹ The available experimental data² seemed to indicate a deficit of events in one region when compared to the Standard Model (SM) prediction.

The physical description of the process can be split into two pieces: the inner bremsstrahlung (IB) contribution and the structure-dependent (SD) contribution, described by two form factors.

Still we lack a complete analysis of the process including q^2 dependence in the form factors, and this is the main purpose of this work. We will use the resonance chiral theory (R χ T) for the description of the form factors.³

Within the SM the SD contributions can be parametrized by a vector form factor and a axial-vector form factor. A tensor form factor would interfere destructively in the region in which the deficit of events occurs.

The axial-vector form factor has already been calculated.⁴ The VVP Green Function so far has been calculated only with one multiplet of vector

*Talk given at the 5th International Workshop on Chiral Dynamics, Theory and Experiment, 18-22 September (2006), Durham/Chapel Hill, NC, (USA); IFIC/06-43 FTUV/06-1121 report. To appear in the Proceedings.

meson resonances and it is not able to fulfil at the same time all the short distance constraints.⁵ Including a second multiplet we can satisfy all the requirements and we obtain a reliable vector form factor:³

$$F_V(q^2) = -\frac{m_{\pi^+}}{3\sqrt{2}B_0FM_{V_1}^2M_{V_2}^2}\frac{c_{000}+c_{010}q^2}{(M_{V_1}^2-q^2)(M_{V_2}^2-q^2)} \quad (1)$$

where c_{000} is fixed by the anomaly and c_{010} is fitted from the $\pi \rightarrow \gamma\gamma^*$ phenomenology.

Finally with this framework we calculate the tensor hadronic matrix element, usually expressed through the so called magnetic susceptibility χ and the vector meson couplings to vector and tensor currents:

$$\chi = -\frac{2}{M_V^2} \simeq -3.37 \text{ GeV}^{-2}, \quad \frac{f_V^\perp}{f_V} = \frac{\langle \bar{q}q \rangle}{M_VF_V^2} \simeq 0.78(1). \quad (2)$$

With the new but preliminary data from the PIBETA collaboration⁶ the mismatch now is gone, as can be seen in tab. 1. Nevertheless we fit the

$E_{e^+}^{\min}(\text{MeV})$	$E_\gamma^{\min}(\text{MeV})$	$\theta_{e\gamma}^{\min}$	$R_{\text{exp}}(\times 10^{-8})^6$	R_{the}^3
50	50	—	2.655 ± 0.058	2.55(9)
10	50	40°	14.59 ± 0.60	14.66(1)
50	10	40°	37.95 ± 0.28	37.70(98)

value for the NP tensor coupling and we get:

$$f_T = (4 \pm 6) \times 10^{-4} \quad (3)$$

which is compatible with zero and with the value dictated by SUSY.¹

Acknowledgements

Work supported by EU FLAVIANet (MRTN-CT-2006-035482), the Spanish MEC (FPA2004-00996), Generalitat Valenciana (GRUPOS03/013 and GV05-164) and by ERDF funds from the EU Commission.

References

1. V. M. Belyaev and I. I. Kogan, Phys. Lett. B **280** (1992) 238.
2. A. A. Poblaguev, Phys. Lett. B **238** (1990) 108;
E. Frlez *et al.*, Phys. Rev. Lett. **93**, 181804 (2004) [arXiv:hep-ex/0312029].
3. V. Mateu and J. Portolés, work in preparation
4. V. Cirigliano, G. Ecker, M. Eidemuller, A. Pich and J. Portoles, Phys. Lett. B **596** (2004) 96

5. M. Knecht and A. Nyffeler, Eur. Phys. J. C **21** (2001) 659;
P. D. Ruiz-Femenia, A. Pich and J. Portolés, JHEP **0307** (2003) 003
6. See D. Poganic proceedings of this workshop